

POSITIVITY OF THE $\bar{\partial}$ -NEUMANN LAPLACIAN

SIQI FU

Dedicated to Professor Linda Rothschild

ABSTRACT. We study the $\bar{\partial}$ -Neumann Laplacian from spectral theoretic perspectives. In particular, we show how pseudoconvexity of a bounded domain is characterized by positivity of the $\bar{\partial}$ -Neumann Laplacian.

MATHEMATICS SUBJECT CLASSIFICATION (2000): 32W05.

KEYWORDS: Pseudoconvex, $\bar{\partial}$ -Neumann Laplacian, Dolbeault cohomology, L^2 -cohomology.

1. INTRODUCTION

Whether or not a given system has positive ground state energy is a widely studied problem with significant repercussions in physics, particularly in quantum mechanics. It follows from the classical Hardy inequality that the bottom of the spectrum of the Dirichlet Laplacian on a domain in $\mathbb{R}^n$ that satisfies the outer cone condition is positive if and only if its inradius is finite (see [D95]). Whereas spectral behavior of the Dirichlet Laplacian is insensitive to boundary geometry, the story for the $\bar{\partial}$ -Neumann Laplacian is different. Since the work of Kohn [Ko63, Ko64] and Hörmander [H65], it has been known that existence and regularity of the $\bar{\partial}$ -Neumann Laplacian closely depend on the underlying geometry (see the surveys [BSt99, Ch99, DK99, FS01] and the monographs [CS99, St09]).

Let Ω be a domain in $\mathbb{C}^n$. It follows from the classical Theorem B of Cartan that if Ω is pseudoconvex, then the Dolbeault cohomology groups $H^{0,q}(\Omega)$ vanish for all $q \geq 1$. (More generally, for any coherent analytic sheaf $\mathcal{F}$ over a Stein manifold, the sheaf cohomology groups $H^q(X, \mathcal{F})$ vanish for all $q \geq 1$.) The converse is also true ([Se53], p. 65). Cartan's Theorem B and its converse were generalized by Laufer [L66] and Siu [Siu67] to a Riemann domain over a Stein manifold. When Ω is bounded, it follows from Hörmander's L^2 -existence theorem for the $\bar{\partial}$ -operator that if Ω is in addition pseudoconvex, then the L^2 -cohomology groups $\tilde{H}^{0,q}(\Omega)$ vanish for $q \geq 1$. The converse of Hörmander's theorem also holds, under the assumption that the interior of the closure of Ω is the domain itself. Sheaf theoretic arguments for the Dolbeault cohomology groups can be modified to give a proof of this fact (cf. [Se53, L66, Siu67, Br83, O88]; see also [Fu05] and Section 3 below).

In this expository paper, we study positivity of the $\bar{\partial}$ -Neumann Laplacian, in connection with the above-mentioned classical results, through the lens of spectral theory. Our emphasis is on the interplay between spectral behavior of the $\bar{\partial}$ -Neumann Laplacian and the geometry of the domains. This is evidently motivated by Marc Kac's famous question "Can one hear the shape of a drum?" [Ka66]. Here we are interested in determining the geometry of a domain in $\mathbb{C}^n$ from the spectrum of the $\bar{\partial}$ -Neumann Laplacian. (See [Fu05, Fu08] for related results.) We make an effort to present a more accessible and self-contained

The author was supported in part by NSF grant DMS-0805852.

treatment, using extensively spectral theoretic language but bypassing sheaf cohomology theory.

2. PRELIMINARIES

In this section, we review the spectral theoretic setup for the $\bar{\partial}$ -Neumann Laplacian. The emphasis here is slightly different from the one in the extant literature (cf. [FK72, CS99]). The $\bar{\partial}$ -Neumann Laplacian is defined through its associated quadratic form. As such, the self-adjoint property and the domain of its square root come out directly from the definition.

Let Q be a non-negative, densely defined, and closed sesquilinear form on a complex Hilbert space H with domain $\mathcal{D}(Q)$. Then Q uniquely determines a non-negative and self-adjoint operator S such that $\mathcal{D}(S^{1/2}) = \mathcal{D}(Q)$ and

$$Q(u, v) = \langle S^{1/2}u, S^{1/2}v \rangle$$

for all $u, v \in \mathcal{D}(Q)$. (See Theorem 4.4.2 in [D95], to which we refer the reader for the necessary spectral theoretic background used in this paper.) For any subspace $L \subset \mathcal{D}(Q)$, let $\lambda(L) = \sup\{Q(u, u) \mid u \in L, \|u\| = 1\}$. For any positive integer j , let

$$(2.1) \quad \lambda_j(Q) = \inf\{\lambda(L) \mid L \subset \mathcal{D}(Q), \dim(L) = j\}.$$

The resolvent set $\rho(S)$ of S consists of all $\lambda \in \mathbb{C}$ such that the operator $S - \lambda I: \mathcal{D}(S) \rightarrow H$ is both one-to-one and onto (and hence has a bounded inverse by the closed graph theorem). The spectrum $\sigma(S)$, the complement of $\rho(S)$ in $\mathbb{C}$, is a non-empty closed subset of $[0, \infty)$. Its bottom $\inf \sigma(S)$ is given by $\lambda_1(Q)$. The essential spectrum $\sigma_e(S)$ is a closed subset of $\sigma(S)$ that consists of isolated eigenvalues of infinite multiplicity and accumulation points of the spectrum. It is empty if and only if $\lambda_j(Q) \rightarrow \infty$ as $j \rightarrow \infty$. In this case, $\lambda_j(Q)$ is the j^{th} eigenvalue of S , arranged in increasing order and repeated according to multiplicity. The bottom of the essential spectrum $\inf \sigma_e(T)$ is the limit of $\lambda_j(Q)$ as $j \rightarrow \infty$. (When $\sigma_e(S) = \emptyset$, we set $\inf \sigma_e(S) = \infty$.)

Let $T_k: H_k \rightarrow H_{k+1}$, $k = 1, 2$, be densely defined and closed operators on Hilbert spaces. Assume that $\mathcal{R}(T_1) \subset \mathcal{N}(T_2)$, where $\mathcal{R}$ and $\mathcal{N}$ denote the range and kernel of the operators. Let T_k^* be the Hilbert space adjoint of T_k , defined in the sense of Von Neumann by

$$\mathcal{D}(T_k^*) = \{u \in H_{k+1} \mid \exists C > 0, |\langle u, T_k v \rangle| \leq C\|v\|, \forall v \in \mathcal{D}(T_k)\}$$

and

$$\langle T_k^* u, v \rangle = \langle u, T_k v \rangle, \quad \text{for all } u \in \mathcal{D}(T_k^*) \text{ and } v \in \mathcal{D}(T_k).$$

Then T_k^* is also densely defined and closed. Let

$$Q(u, v) = \langle T_1^* u, T_1^* v \rangle + \langle T_2 u, T_2 v \rangle$$

with its domain given by $\mathcal{D}(Q) = \mathcal{D}(T_1^*) \cap \mathcal{D}(T_2)$. The following proposition elucidates the above approach to the $\bar{\partial}$ -Neumann Laplacian.

Proposition 2.1. *$Q(u, v)$ is a densely defined, closed, non-negative sesquilinear form. The associated self-adjoint operator $\square$ is given by*

$$(2.2) \quad \mathcal{D}(\square) = \{f \in H_2 \mid f \in \mathcal{D}(Q), T_2 f \in \mathcal{D}(T_2^*), T_1^* f \in \mathcal{D}(T_1)\}, \quad \square = T_1 T_1^* + T_2^* T_2.$$

Proof. The closedness of Q follows easily from that of T_1 and T_2 . The non-negativity is evident. We now prove that $\mathcal{D}(Q)$ is dense in H_2 . Since $\mathcal{N}(T_2)^\perp = \overline{\mathcal{R}(T_2^*)} \subset \mathcal{N}(T_1^*)$ and

$$\mathcal{D}(T_2) = \mathcal{N}(T_2) \oplus (\mathcal{D}(T_2) \cap \mathcal{N}(T_2)^\perp),$$

we have

$$\mathcal{D}(Q) = \mathcal{D}(T_1^*) \cap \mathcal{D}(T_2) = (\mathcal{N}(T_2) \cap \mathcal{D}(T_1^*)) \oplus (\mathcal{D}(T_2) \cap \mathcal{N}(T_2)^\perp).$$

Since $\mathcal{D}(T_1^*)$ and $\mathcal{D}(T_2)$ are dense in H_2 , $\mathcal{D}(Q)$ is dense in $\mathcal{N}(T_2) \oplus \mathcal{N}(T_2)^\perp = H_2$.

It follows from the above definition of $\square$ that $f \in \mathcal{D}(\square)$ if and only if $f \in \mathcal{D}(Q)$ and there exists a $g \in H_2$ such that

$$(2.3) \quad Q(u, f) = \langle u, g \rangle, \quad \text{for all } u \in \mathcal{D}(Q)$$

(cf. Lemma 4.4.1 in [D95]). Thus

$$\mathcal{D}(\square) \supset \{f \in H_2 \mid f \in \mathcal{D}(Q), T_2 f \in \mathcal{D}(T_2^*), T_1^* f \in \mathcal{D}(T_1)\}.$$

We now prove the opposite containment. Suppose $f \in \mathcal{D}(\square)$. For any $u \in \mathcal{D}(T_2)$, we write $u = u_1 + u_2 \in (\mathcal{N}(T_1^*) \cap \mathcal{D}(T_2)) \oplus \mathcal{N}(T_1^*)^\perp$. Note that $\mathcal{N}(T_1^*)^\perp \subset \mathcal{R}(T_2^*)^\perp = \mathcal{N}(T_2)$. It follows from (2.3) that

$$|\langle T_2 u, T_2 f \rangle| = |\langle T_2 u_1, T_2 f \rangle| = |Q(u_1, f)| = |\langle u_1, g \rangle| \leq \|u\| \cdot \|g\|.$$

Hence $T_2 f \in \mathcal{D}(T_2^*)$. The proof of $T_1^* f \in \mathcal{D}(T_1)$ is similar. For any $w \in \mathcal{D}(T_1^*)$, we write $w = w_1 + w_2 \in (\mathcal{N}(T_2) \cap \mathcal{D}(T_1^*)) \oplus \mathcal{N}(T_2)^\perp$. Note that $\mathcal{N}(T_2)^\perp = \overline{\mathcal{R}(T_2^*)} \subset \mathcal{N}(T_1^*)$. Therefore, by (2.3),

$$|\langle T_1^* w, T_1^* f \rangle| = |\langle T_1^* w_1, T_1^* f \rangle| = |Q(w_1, f)| = |\langle w_1, g \rangle| \leq \|w\| \cdot \|g\|.$$

Hence $T_1^* f \in \mathcal{D}(T_1^{**}) = \mathcal{D}(T_1)$. It follows from the definition of $\square$ that for any $f \in \mathcal{D}(\square)$ and $u \in \mathcal{D}(Q)$,

$$\langle \square f, u \rangle = \langle \square^{1/2} f, \square^{1/2} u \rangle = Q(f, u) = \langle T_1^* f, T_1^* u \rangle + \langle T_2 f, T_2 u \rangle = \langle (T_1 T_1^* + T_2^* T_2) f, u \rangle.$$

Hence $\square = T_1 T_1^* + T_2^* T_2$. $\square$

The following proposition is well-known (compare [H65], Theorem 1.1.2 and Theorem 1.1.4; [C83], Proposition 3; and [Sh92], Proposition 2.3). We provide a proof here for completeness.

Proposition 2.2. *$\inf \sigma(\square) > 0$ if and only if $\mathcal{R}(T_1) = \mathcal{N}(T_2)$ and $\mathcal{R}(T_2)$ is closed.*

Proof. Assume $\inf \sigma(\square) > 0$. Then 0 is in the resolvent set of $\square$ and hence $\square$ has a bounded inverse $G: H_2 \rightarrow \mathcal{D}(\square)$. For any $u \in H_2$, write $u = T_1 T_1^* G u + T_2^* T_2 G u$. If $u \in \mathcal{N}(T_2)$, then $0 = (T_2 u, T_2 G u) = (T_2 T_2^* T_2 G u, T_2 G u) = (T_2^* T_2 G u, T_2^* T_2 G u)$. Hence $T_2^* T_2 G u = 0$ and $u = T_1 T_1^* G u$. Therefore, $\mathcal{R}(T_1) = \mathcal{N}(T_2)$. Similarly, $\mathcal{R}(T_2^*) = \mathcal{N}(T_1^*)$. Therefore T_2^* and hence T_2 have closed range. To prove the opposite implication, we write $u = u_1 + u_2 \in \mathcal{N}(T_2) \oplus \mathcal{N}(T_2)^\perp$, for any $u \in \mathcal{D}(Q)$. Note that $u_1, u_2 \in \mathcal{D}(Q)$. It follows from $\mathcal{N}(T_2) = \mathcal{R}(T_1)$ and the closed range property of T_2 that there exists a positive constant c such that $c\|u_1\|^2 \leq \|T_1^* u_1\|^2$ and $c\|u_2\|^2 \leq \|T_2 u_2\|^2$. Thus

$$c\|u\|^2 = c(\|u_1\|^2 + \|u_2\|^2) \leq \|T_1^* u_1\|^2 + \|T_2 u_2\|^2 = Q(u, u).$$

Hence $\inf \sigma(\square) \geq c > 0$ (cf. Theorem 4.3.1 in [D95]). $\square$

Let $\mathcal{N}(Q) = \mathcal{N}(T_1^*) \cap \mathcal{N}(T_2)$. Note that when it is non-trivial, $\mathcal{N}(Q)$ is the eigenspace of the zero eigenvalue of $\square$. When $\mathcal{R}(T_1)$ is closed, $\mathcal{N}(T_2) = \mathcal{R}(T_1) \oplus \mathcal{N}(Q)$. For a subspace $L \subset H_2$, denote by $P_{L^\perp}$ the orthogonal projection onto $L^\perp$ and $T_2|_{L^\perp}$ the restriction of T_2 to $L^\perp$. The next proposition clarifies and strengthens the second part of Lemma 2.1 in [Fu05].

Proposition 2.3. *The following statements are equivalent:*

- (1) $\inf \sigma_e(\square) > 0$.
- (2) $\mathcal{R}(T_1)$ and $\mathcal{R}(T_2)$ are closed and $\mathcal{N}(Q)$ is finite dimensional.
- (3) There exists a finite dimensional subspace $L \subset \mathcal{D}(T_1^*) \cap \mathcal{N}(T_2)$ such that $\mathcal{N}(T_2) \cap L^\perp = P_{L^\perp}(\mathcal{R}(T_1))$ and $\mathcal{R}(T_2|_{L^\perp})$ is closed.

Proof. We first prove (1) implies (2). Suppose $a = \inf \sigma_e(\square) > 0$. If $\inf \sigma(\square) > 0$, then $\mathcal{N}(Q)$ is trivial and (2) follows from Proposition 2.2. Suppose $\inf \sigma(\square) = 0$. Then $\sigma(\square) \cap [0, a)$ consists only of isolated points, all of which are eigenvalues of finite multiplicity of $\square$ (cf. Theorem 4.5.2 in [D95]). Hence $\mathcal{N}(Q)$, the eigenspace of the eigenvalue 0, is finite dimensional. Choose a sufficiently small $c > 0$ so that $\sigma(\square) \cap [0, c) = \{0\}$. By the spectral theorem for self-adjoint operators (cf. Theorem 2.5.1 in [D95]), there exists a finite regular Borel measure μ on $\sigma(\square) \times \mathbb{N}$ and a unitary transformation $U: H_2 \rightarrow L^2(\sigma(\square) \times \mathbb{N}, d\mu)$ such that $U\square U^{-1} = M_x$, where $M_x\varphi(x, n) = x\varphi(x, n)$ is the multiplication operator by x on $L^2(\sigma(\square) \times \mathbb{N}, d\mu)$. Let $P_{\mathcal{N}(Q)}$ be the orthogonal projection onto $\mathcal{N}(Q)$. For any $f \in \mathcal{D}(Q) \cap \mathcal{N}(Q)^\perp$,

$$UP_{\mathcal{N}(Q)}f = \chi_{[0, c)}Uf = 0,$$

where $\chi_{[0, c)}$ is the characteristic function of $[0, c)$. Hence Uf is supported on $[c, \infty)$. Therefore,

$$Q(f, f) = \int_{\sigma(\square) \times \mathbb{N}} x|Uf|^2 d\mu \geq c\|Uf\|^2 = c\|f\|^2.$$

It then follows from Theorem 1.1.2 in [H65] that both T_1 and T_2 have closed range.

To prove (2) implies (1), we use Theorem 1.1.2 in [H65] in the opposite direction: There exists a positive constant c such that

$$(2.4) \quad c\|f\|^2 \leq Q(f, f), \quad \text{for all } f \in \mathcal{D}(Q) \cap \mathcal{N}(Q)^\perp.$$

Proving by contradiction, we assume $\inf \sigma_e(\square) = 0$. Let ε be any positive number less than c . Since $L_{[0, \varepsilon)} = \mathcal{R}(\chi_{[0, \varepsilon)}(\square))$ is infinite dimensional (cf. Lemma 4.1.4 in [D95]), there exists a non-zero $g \in L_{[0, \varepsilon)}$ such that $g \perp \mathcal{N}(Q)$. However,

$$Q(g, g) = \int_{\sigma(\square) \times \mathbb{N}} x\chi_{[0, \varepsilon)}|Ug|^2 d\mu \leq \varepsilon\|Ug\|^2 = \varepsilon\|g\|^2,$$

contradicting (2.4).

We do some preparations before proving the equivalence of (3) with (1) and (2). Let L be any finite dimensional subspace of $\mathcal{D}(T_1^*) \cap \mathcal{N}(T_2)$. Let $H'_2 = H_2 \ominus L$. Let $T'_2 = T_2|_{H'_2}$ and let $T'^*_1 = T^*_1|_{H'_2}$. Then $T'_2: H'_2 \rightarrow H_3$ and $T'^*_1: H'_2 \rightarrow H_1$ are densely defined, closed operators. Let $T'_1: H_1 \rightarrow H'_2$ be the adjoint of T'^*_1 . It follows from the definitions that $\mathcal{D}(T_1) \subset \mathcal{D}(T'_1)$. The finite dimensionality of L implies the opposite containment. Thus, $\mathcal{D}(T_1) = \mathcal{D}(T'_1)$. For any $f \in \mathcal{D}(T_1)$ and $g \in \mathcal{D}(T'^*_1) = \mathcal{D}(T^*_1) \cap L^\perp$,

$$\langle T'_1 f, g \rangle = \langle f, T'^*_1 g \rangle = \langle f, T^*_1 g \rangle = \langle T_1 f, g \rangle.$$

Hence $T'_1 = P_{L^\perp} \circ T_1$ and $\mathcal{R}(T'_1) = P_{L^\perp}(\mathcal{R}(T_1)) \subset \mathcal{N}(T'_2)$. Let

$$Q'(f, g) = \langle T'^*_1 f, T'^*_1 g \rangle + \langle T'_2 f, T'_2 g \rangle$$

be the associated sesquilinear form on H'_2 with $\mathcal{D}(Q') = \mathcal{D}(Q) \cap L^\perp$.

We are now in position to prove that (2) implies (3). In this case, we can take $L = \mathcal{N}(Q)$. By Theorem 1.1.2 in [H65], there exists a positive constant c such that

$$Q(f, f) = Q'(f, f) \geq c\|f\|^2, \quad \text{for all } f \in \mathcal{D}(Q').$$

We then obtain (3) by applying Proposition 2.2 to T'_1 , T'_2 , and $Q'(f, g)$.

Finally, we prove (3) implies (1). Applying Proposition 2.2 in the opposite direction, we know that there exists a positive constant c such that

$$Q(f, f) \geq c\|f\|^2, \quad \text{for all } f \in \mathcal{D}(Q) \cap L^\perp.$$

The rest of the proof follows the same lines of the above proof of the implication (2) $\Rightarrow$ (1), with $\mathcal{N}(Q)$ there replaced by L . $\square$

We now recall the definition of the $\bar{\partial}$ -Neumann Laplacian on a complex manifold. Let X be a complex hermitian manifold of dimension n . Let $C_{(0,q)}^\infty(X) = C^\infty(X, \Lambda^{0,q}T^*X)$ be the space of smooth $(0, q)$ -forms on X . Let $\bar{\partial}_q: C_{(0,q)}^\infty(X) \rightarrow C_{(0,q+1)}^\infty(X)$ be the composition of the exterior differential operator and the projection onto $C_{(0,q+1)}^\infty(X)$.

Let Ω be a domain in X . For $u, v \in C_{(0,q)}^\infty(X)$, let $\langle u, v \rangle$ be the point-wise inner product of u and v , and let

$$\langle\langle u, v \rangle\rangle_\Omega = \int_\Omega \langle u, v \rangle dV$$

be the inner product of u and v over Ω . Let $L_{(0,q)}^2(\Omega)$ be the completion of the space of compactly supported forms in $C_{(0,q)}^\infty(\Omega)$ with respect to the above inner product. The operator $\bar{\partial}_q$ has a closed extension on $L_{(0,q)}^2(\Omega)$. We also denote the closure by $\bar{\partial}_q$. Thus $\bar{\partial}_q: L_{(0,q)}^2(\Omega) \rightarrow L_{(0,q+1)}^2(\Omega)$ is densely defined and closed. Let $\bar{\partial}_q^*$ be its adjoint. For $1 \leq q \leq n-1$, let

$$Q_q(u, v) = \langle\langle \bar{\partial}_q u, \bar{\partial}_q v \rangle\rangle_\Omega + \langle\langle \bar{\partial}_{q-1}^* u, \bar{\partial}_{q-1}^* v \rangle\rangle_\Omega$$

be the sesquilinear form on $L_{(0,q)}^2(\Omega)$ with domain $\mathcal{D}(Q_q) = \mathcal{D}(\bar{\partial}_q) \cap \mathcal{D}(\bar{\partial}_{q-1}^*)$. The self-adjoint operator $\square_q$ associated with Q_q is called the $\bar{\partial}$ -Neumann Laplacian on $L_{(0,q)}^2(\Omega)$. It is an elliptic operator with non-coercive boundary conditions [KN65].

The Dolbeault and L^2 -cohomology groups on Ω are defined respectively by

$$H^{0,q}(\Omega) = \frac{\{f \in C_{(0,q)}^\infty(\Omega) \mid \bar{\partial}_q f = 0\}}{\{\bar{\partial}_{q-1} g \mid g \in C_{(0,q-1)}^\infty(\Omega)\}} \quad \text{and} \quad \tilde{H}^{0,q}(\Omega) = \frac{\{f \in L_{(0,q)}^2(\Omega) \mid \bar{\partial}_q f = 0\}}{\{\bar{\partial}_{q-1} g \mid g \in L_{(0,q-1)}^2(\Omega)\}}.$$

These cohomology groups are in general not isomorphic. For example, when a complex variety is deleted from Ω , the L^2 -cohomology group remains the same but the Dolbeault cohomology group could change from trivial to infinite dimensional. As noted in the paragraph preceding Proposition 2.3, when $\mathcal{R}(\bar{\partial}_{q-1})$ is closed in $L_{(0,q)}^2(\Omega)$, $\tilde{H}^{0,q}(\Omega) \cong \mathcal{N}(\square_q)$. We refer the reader to [De] for an extensive treatise on the subject and to [H65] and [O82] for results relating these cohomology groups.

3. POSITIVITY OF THE SPECTRUM AND ESSENTIAL SPECTRUM

Laufer proved in [L75] that for any open subset of a Stein manifold, if a Dolbeault cohomology group is finite dimensional, then it is trivial. In this section, we establish the following L^2 -analogue of this result on a bounded domain in a Stein manifold:

Theorem 3.1. *Let $\Omega \subset\subset X$ be a domain in a Stein manifold X with C^1 boundary. Let $\square_q$, $1 \leq q \leq n-1$, be the $\bar{\partial}$ -Neumann Laplacian on $L_{(0,q)}^2(\Omega)$. Assume that $\mathcal{N}(\square_q) \subset W^1(\Omega)$. Then $\inf \sigma(\square_q) > 0$ if and only if $\inf \sigma_e(\square_q) > 0$.*

The proof of Theorem 3.1 follows the same line of arguments as Laufer's. We provide the details below.

Let $H^\infty(\Omega)$ be the space of bounded holomorphic functions on Ω . For any $f \in H^\infty(\Omega)$, let M_f be the multiplication operator by f :

$$M_f: L^2_{(0,q)}(\Omega) \rightarrow L^2_{(0,q)}(\Omega), \quad M_f(u) = fu.$$

Then M_f induces an endomorphism on $\tilde{H}^{0,q}(\Omega)$. Let $\mathcal{I}$ be set of all holomorphic functions $f \in H^\infty(\Omega)$ such that $M_f = 0$ on $\tilde{H}^{0,q}(\Omega)$. Evidently, $\mathcal{I}$ is an ideal of $H^\infty(\Omega)$. Assume $\inf \sigma_e(\square_q) > 0$. To show that $\tilde{H}^{0,q}(\Omega)$ is trivial, it suffices to show that $1 \in \mathcal{I}$.

Lemma 3.2. *Let ξ be a holomorphic vector field on X and let $f \in \mathcal{I}$. Then $\xi(f) \in \mathcal{I}$.*

Proof. Let $D = \xi \lrcorner \partial: C^\infty_{(0,q)}(\Omega) \rightarrow C^\infty_{(0,q)}(\Omega)$, where $\lrcorner$ denotes the contraction operator. It is easy to check that D commute with the $\bar{\partial}$ operator. Therefore, D induces an endomorphism on $\tilde{H}^{0,q}(\Omega)$. (Recall that under the assumption, $\tilde{H}^{0,q}(\Omega) \cong \mathcal{N}(\square_q) \subset W^1(\Omega)$.) For any $u \in \mathcal{N}(\square_q)$,

$$D(fu) - fD(u) = \xi \lrcorner \partial(fu) - f\xi \lrcorner \partial u = \xi(f)u.$$

Notice that Ω is locally starlike near the boundary. Using partition of unity and the Friedrichs Lemma, we obtain $[D(fu)] = 0$. Therefore, $[\xi(f)u] = [D(fu)] - [fD(u)] = [0]$. $\square$

We now return to the proof of the theorem. Let $F = (f_1, \dots, f_{n+1}): X \rightarrow \mathbb{C}^{2n+1}$ be a proper embedding of X into $\mathbb{C}^{2n+1}$ (cf. Theorem 5.3.9 in [H91]). Since Ω is relatively compact in X , $f_j \in H^\infty(\Omega)$. For any f_j , let $P_j(\lambda)$ be the characteristic polynomial of $M_{f_j}: \tilde{H}^{0,q}(\Omega) \rightarrow \tilde{H}^{0,q}(\Omega)$. By the Cayley-Hamilton theorem, $P_j(M_{f_j}) = 0$ (cf. Theorem 2.4.2 in [HJ85]). Thus $P_j(f_j) \in \mathcal{I}$.

The number of points in the set $\{(\lambda_1, \lambda_2, \dots, \lambda_{2n+1}) \in \mathbb{C}^{2n+1} \mid P_j(\lambda_j) = 0, 1 \leq j \leq 2n+1\}$ is finite. Since $F: X \rightarrow \mathbb{C}^{2n+1}$ is one-to-one, the number of common zeroes of $P_j(f_j(z))$, $1 \leq j \leq 2n+1$, on X is also finite. Denote these zeroes by z^k , $1 \leq k \leq N$. For each z^k , let g_k be a function in $\mathcal{I}$ whose vanishing order at z^k is minimal. (Since $P_j(f_j) \in \mathcal{I}$, $g_k \neq 0$.) We claim that $g_k(z^k) \neq 0$. Suppose otherwise $g_k(z^k) = 0$. Since there exists a holomorphic vector field ξ on X with any prescribed holomorphic tangent vector at any given point (cf. Corollary 5.6.3 in [H91]), one can find an appropriate choice of ξ so that $\xi(g_j)$ vanishes to lower order at z^k . According to Lemma 3.2, $\xi(g_j) \in \mathcal{I}$. We thus arrive at a contradiction.

Now we know that there are holomorphic functions, $P_j(f_j)$, $1 \leq j \leq 2n+1$, and g_k , $1 \leq k \leq N$, that have no common zeroes on X . It then follows that there exist holomorphic functions h_j on X such that

$$\sum P_j(f_j)h_j + \sum g_k h_k = 1.$$

(See, for example, Corollary 16 on p. 244 in [GR65], Theorem 7.2.9 in [H91], and Theorem 7.2.5 in [Kr01]. Compare also Theorem 2 in [Sk72].) Since $P_j(f_j) \in \mathcal{I}$, $g_k \in \mathcal{I}$, and $h_j \in H^\infty(\Omega)$, we have $1 \in \mathcal{I}$. We thus conclude the proof of Theorem 3.1.

Remark. (1) Unlike the above-mentioned result of Laufer on the Dolbeault cohomology groups [L75], Theorem 3.1 is not expected to hold if the boundedness condition on Ω is removed (compare [W84]). It would be interesting to know whether Theorem 3.1 remains true if the assumption $\mathcal{N}(\square_q) \subset W^1(\Omega)$ is dropped and whether it remains true for unbounded pseudoconvex domains.

(2) Notice that in the above proof, we use the fact that $\mathcal{R}(\bar{\partial}_{q-1})$ is closed, as a consequence of the assumption $\inf \sigma_e(\square_q) > 0$ by Proposition 2.3. It is well known that for any infinite dimensional Hilbert space H , there exists a subspace R of H such that H/R is finite dimensional but R is not closed. However, the construction of such a subspace usually involves Zorn's lemma (equivalently, the axiom of choice). It would be of interest to know whether there exists a domain Ω in a Stein manifold such that $\tilde{H}^{0,q}(\Omega)$ is finite dimensional but $\mathcal{R}(\bar{\partial}_{q-1})$ is not closed.

(3) We refer the reader to [Sh09] for related results on the relationship between triviality and finite dimensionality of the L^2 -cohomology groups using the $\bar{\partial}$ -Cauchy problem. We also refer the reader to [B02] for a related result on embedded CR manifolds.

4. HEARING PSEUDOCONVEXITY

The following theorem illustrates that one can easily determine pseudoconvexity from the spectrum of the $\bar{\partial}$ -Neumann Laplacian.

Theorem 4.1. *Let Ω be a bounded domain in $\mathbb{C}^n$ such that $\text{int}(\text{cl}(\Omega)) = \Omega$. Then the following statements are equivalent:*

- (1) Ω is pseudoconvex.
- (2) $\inf \sigma(\square_q) > 0$, for all $1 \leq q \leq n-1$.
- (3) $\inf \sigma_e(\square_q) > 0$, for all $1 \leq q \leq n-1$.

The implication (1) $\Rightarrow$ (2) is a consequence of Hörmander's fundamental L^2 -estimates of the $\bar{\partial}$ -operator [H65], in light of Proposition 2.2, and it holds without the assumption $\text{int}(\text{cl}(\Omega)) = \Omega$. The implications (2) $\Rightarrow$ (1) and (3) $\Rightarrow$ (1) are consequences of the sheaf cohomology theory dated back to Oka and Cartan (cf. [Se53, L66, Siu67, Br83, O88]). A elementary proof of (2) implying (1), as explained in [Fu05], is given below. The proof uses sheaf cohomology arguments in [L66]. When adapting Laufer's method to study the L^2 -cohomology groups, one encounters a difficulty: While the restriction to the complex hyperplane of the smooth function resulting from the sheaf cohomology arguments for the Dolbeault cohomology groups is well-defined, the restriction of the corresponding L^2 function is not. This difficulty was overcome in [Fu05] by appropriately modifying the construction of auxiliary $(0, q)$ -forms (see the remark after the proof for more elaborations on this point).

We now show that (2) implies (1). Proving by contradiction, we assume that Ω is not pseudoconvex. Then there exists a domain $\tilde{\Omega} \supsetneq \Omega$ such that every holomorphic function on Ω extends to $\tilde{\Omega}$. Since $\text{int}(\text{cl}(\Omega)) = \Omega$, $\tilde{\Omega} \setminus \text{cl}(\Omega)$ is non-empty. After a translation and a unitary transformation, we may assume that the origin is in $\tilde{\Omega} \setminus \text{cl}(\Omega)$ and there is a point z^0 in the intersection of z_n -plane with Ω that is in the same connected component of the intersection of the z_n -plane with $\tilde{\Omega}$.

Let m be a positive integer (to be specified later). Let $k_q = n$. For any $\{k_1, \dots, k_{q-1}\} \subset \{1, 2, \dots, n-1\}$, we define

$$(4.1) \quad u(k_1, \dots, k_q) = \frac{(q-1)! (\bar{z}_{k_1} \cdots \bar{z}_{k_q})^{m-1}}{r_m^q} \sum_{j=1}^q (-1)^j \bar{z}_{k_j} d\bar{z}_{k_1} \wedge \cdots \wedge \widehat{d\bar{z}_{k_j}} \wedge \cdots \wedge d\bar{z}_{k_q},$$

where $r_m = |z_1|^{2m} + \cdots + |z_n|^{2m}$. As usual, $\widehat{d\bar{z}_{k_j}}$ indicates the deletion of $d\bar{z}_{k_j}$ from the wedge product. Evidently, $u(k_1, \dots, k_q) \in L^2_{(0,q-1)}(\Omega)$ is a smooth form on $\mathbb{C}^n \setminus \{0\}$. Moreover,

$u(k_1, \dots, k_q)$ is skew-symmetric with respect to the indices $(k_1, \dots, k_{q-1})$. In particular, $u(k_1, \dots, k_q) = 0$ when two k_j 's are identical.

We now fix some notional conventions. Let $K = (k_1, \dots, k_q)$ and J a collection of indices from $\{k_1, \dots, k_q\}$. Write $d\bar{z}_K = d\bar{z}_{k_1} \wedge \dots \wedge d\bar{z}_{k_q}$, $\bar{z}_K^{m-1} = (\bar{z}_{k_1} \dots \bar{z}_{k_q})^{m-1}$, and $\widetilde{d\bar{z}_{k_j}} = d\bar{z}_{k_1} \wedge \dots \wedge \widehat{d\bar{z}_{k_j}} \wedge \dots \wedge d\bar{z}_{k_q}$. Denote by $(k_1, \dots, k_q \mid J)$ the tuple of remaining indices after deleting those in J from $(k_1, \dots, k_q)$. For example, $(2, 5, 3, 1 \mid (4, 1, 6 \mid 4, 6)) = (2, 5, 3)$.

It follows from a straightforward calculation that

$$\begin{aligned}
 \bar{\partial}u(k_1, \dots, k_q) &= -\frac{q!m\bar{z}_K^{m-1}}{r_m^{q+1}}(r_m d\bar{z}_K + (\sum_{\ell=1}^n \bar{z}_\ell^{m-1} z_\ell^m d\bar{z}_\ell) \wedge (\sum_{j=1}^q (-1)^j \bar{z}_{k_j} \widetilde{d\bar{z}_{k_j}})) \\
 &= -\frac{q!m\bar{z}_K^{m-1}}{r_m^{q+1}} \sum_{\substack{\ell=1 \\ \ell \neq k_1, \dots, k_q}}^n z_\ell^m \bar{z}_\ell^{m-1} (\bar{z}_\ell d\bar{z}_K + d\bar{z}_\ell \wedge \sum_{j=1}^q (-1)^j \bar{z}_{k_j} \widetilde{d\bar{z}_{k_j}}) \\
 (4.2) \quad &= m \sum_{\ell=1}^{n-1} z_\ell^m u(\ell, k_1, \dots, k_q).
 \end{aligned}$$

It follows that $u(1, \dots, n)$ is a $\bar{\partial}$ -closed $(0, n-1)$ -form.

By Proposition 2.2, we have $\mathcal{R}(\bar{\partial}_{q-1}) = \mathcal{N}(\bar{\partial}_q)$ for all $1 \leq q \leq n-1$. We now solve the $\bar{\partial}$ -equations inductively, using $u(1, \dots, n)$ as initial data. Let $v \in L^2_{(0, n-2)}(\Omega)$ be a solution to $\bar{\partial}v = u(1, \dots, n)$. For any $k_1 \in \{1, \dots, n-1\}$, define

$$w(k_1) = -mz_{k_1}^m v + (-1)^{1+k_1} u(1, \dots, n \mid k_1).$$

Then it follows from (4.2) that $\bar{\partial}w(k_1) = 0$. Let $v(k_1) \in L^2_{(0, n-3)}(\Omega)$ be a solution of $\bar{\partial}v(k_1) = w(k_1)$.

Suppose for any $(q-1)$ -tuple $K' = (k_1, \dots, k_{q-1})$ of integers from $\{1, \dots, n-1\}$, $q \geq 2$, we have constructed $v(K') \in L^2_{(0, n-q-1)}(\Omega)$ such that it is skew-symmetric with respect to the indices and satisfies

$$(4.3) \quad \bar{\partial}v(K') = m \sum_{j=1}^{q-1} (-1)^j z_{k_j}^m v(K' \mid k_j) + (-1)^{q+|K'|} u(1, \dots, n \mid K')$$

where $|K'| = k_1 + \dots + k_{q-1}$ as usual. We now construct a $(0, n-q-2)$ -forms $v(K)$ satisfying (4.3) for any q -tuple $K = (k_1, \dots, k_q)$ of integers from $\{1, \dots, n-1\}$ (with K' replaced by K). Let

$$w(K) = m \sum_{j=1}^q (-1)^j z_{k_j}^m v(K \mid k_j) + (-1)^{q+|K|} u(1, \dots, n \mid K).$$

Then it follows from (4.2) that

$$\begin{aligned}
\bar{\partial}w(K) &= m \sum_{j=1}^q (-1)^j z_{k_j}^m \bar{\partial}v(K | k_j) + (-1)^{q+|K|} \bar{\partial}u(1, \dots, n | K) \\
&= m \sum_{j=1}^q (-1)^j z_{k_j}^m \left(m \sum_{1 \leq i < j} (-1)^i z_{k_i}^m v(K | k_j, k_i) + m \sum_{j < i \leq q} (-1)^{i-1} z_{k_i}^m v(K | k_j, k_i) \right. \\
&\quad \left. - (-1)^{q+|K|-k_j} u(1, \dots, n | (K | k_j)) \right) + (-1)^{q+|K|} \bar{\partial}u(1, \dots, n | K) \\
&= (-1)^{q+|K|} \left(-m \sum_{j=1}^q (-1)^{j-k_j} z_{k_j}^m u(1, \dots, n | (K | k_j)) + \bar{\partial}u(1, \dots, n | K) \right) \\
&= (-1)^{q+|K|} \left(-m \sum_{j=1}^q z_{k_j}^m u(k_j, (1, \dots, n | K)) + \bar{\partial}u(1, \dots, n | K) \right) = 0
\end{aligned}$$

Therefore, by the hypothesis, there exists a $v(K) \in L^2_{(0, n-q-2)}(\Omega)$ such that $\bar{\partial}v(K) = w(K)$. Since $w(K)$ is skew-symmetric with respect to indices K , we may also choose a likewise $v(K)$. This then concludes the inductive step.

Now let

$$F = w(1, \dots, n-1) = m \sum_{j=1}^{n-1} z_j^m v(1, \dots, j, \dots, n-1) - (-1)^{n+\frac{n(n-1)}{2}} u(n),$$

where $u(n) = -\bar{z}_n^m / r_m$, as given by (4.1). Then $F(z) \in L^2(\Omega)$ and $\bar{\partial}F(z) = 0$. By the hypothesis, $F(z)$ has a holomorphic extension to $\tilde{\Omega}$. We now restrict $F(z)$ to the coordinate hyperplane $z' = (z_1, \dots, z_{n-1}) = 0$. Notice that so far we only choose the $v(K)$'s and $w(K)$'s from L^2 -spaces. The restriction to the coordinate hyperplane $z' = 0$ is not well-defined. To overcome this difficulty, we choose $m > 2(n-1)$. For sufficiently small $\varepsilon > 0$ and $\delta > 0$,

$$\begin{aligned}
&\left\{ \int_{\{|z'| < \varepsilon\} \cap \Omega} |(F + (-1)^{n+\frac{n(n-1)}{2}} u(n))(\delta z', z_n)|^2 dV(z) \right\}^{1/2} \\
&\leq m \delta^m \varepsilon^m \sum_{j=1}^{n-1} \left\{ \int_{\{|z'| < \varepsilon\} \cap \Omega} |v(1, \dots, \hat{j}, \dots, n-1)(\delta z', z_n)|^2 dV(z) \right\}^{1/2} \\
&\leq m \delta^{m-2(n-1)} \varepsilon^m \sum_{j=1}^{n-1} \|v(1, \dots, \hat{j}, \dots, n-1)\|_{L^2(\Omega)}.
\end{aligned}$$

Letting $\delta \rightarrow 0$, we then obtain

$$F(0, z_n) = -(-1)^{n+\frac{n(n-1)}{2}} u(n)(0, z_n) = (-1)^{n+\frac{n(n-1)}{2}} z_n^{-m}.$$

for z_n near z_n^0 . (Recall that $z^0 \in \Omega$ is in the same connected component of $\{z' = 0\} \cap \tilde{\Omega}$ as the origin.) This contradicts the analyticity of F near the origin. We therefore conclude the proof of Theorem 4.1.

Remark. (1) The above proof of the implication (2) $\Rightarrow$ (1) uses only the fact that the L^2 -cohomology groups $\tilde{H}^{0,q}(\Omega)$ are trivial for all $1 \leq q \leq n-1$. Under the (possibly) stronger assumption $\inf \sigma(\square_q) > 0$, $1 \leq q \leq n-1$, the difficulty regarding the restriction of the

L^2 function to the complex hyperplane in the proof becomes superficial. In this case, the $\bar{\partial}$ -Neumann Laplacian $\square_q$ has a bounded inverse. The interior ellipticity of the $\bar{\partial}$ -complex implies that one can in fact choose the forms $v(K)$ and $w(K)$ to be smooth inside Ω , using the canonical solution operator to the $\bar{\partial}$ -equation. Therefore, in this case, the restriction to $\{z' = 0\} \cap \Omega$ is well-defined. Hence one can choose $m = 1$. This was indeed the choice in [L66], where the forms involved are smooth and the restriction poses no problem. It is interesting to note that by having the freedom to choose m sufficiently large, one can leave out the use of interior ellipticity. Also, the freedom to choose m becomes crucial when one proves an analogue of Theorem 4.1 for the Kohn Laplacian because the $\bar{\partial}_b$ -complex is no longer elliptic. The construction of $u(k_1, \dots, k_q)$ in (4.1) with the exponent m was introduced in [Fu05] to handle this difficulty.

(2) One can similarly give a proof of the implication (3) $\Rightarrow$ (1). Indeed, the above proof can be easily modified to show that the finite dimensionality of $\tilde{H}^{0,q}(\Omega)$, $1 \leq q \leq n-1$, implies the pseudoconvexity of Ω . In this case, the $u(K)$'s are defined by

$$u(k_1, \dots, k_q) = \frac{(\alpha + q - 1)! \bar{z}_n^{m\alpha} (\bar{z}_{k_1} \cdots \bar{z}_{k_q})^{m-1}}{r_m^{\alpha+q}} \sum_{j=1}^q (-1)^j \bar{z}_{k_j} d\bar{z}_{k_1} \wedge \cdots \wedge \widehat{d\bar{z}_{k_j}} \wedge \cdots \wedge d\bar{z}_{k_q},$$

where α is any non-negative integers. One now fixes a choice of $m > 2(n-1)$ and let α runs from 0 to N for a sufficiently large N , depending on the dimensions of the L^2 -cohomology groups. We refer the reader to [Fu05] for details.

(3) As noted in Sections 2 and 3, unlike the Dolbeault cohomology case, one cannot remove the assumption $\text{int}(\text{cl}(\Omega)) = \Omega$ or the boundedness condition on Ω from Theorem 4.1. For example, a bounded pseudoconvex domain in $\mathbb{C}^n$ with a complex analytic variety removed still satisfies condition (2) in Theorem 3.1.

(4) As in [L66], Theorem 4.1 remains true for a Stein manifold. More generally, as a consequence of Andreotti-Grauert's theory [AG62], the q -convexity of a bounded domain Ω in a Stein manifold such that $\text{int}(\text{cl}(\Omega)) = \Omega$ is characterized by $\inf \sigma(\square_k) > 0$ or $\inf \sigma_e(\square_k) > 0$ for all $q \leq k \leq n-1$.

(5) It follows from Theorem 3.1 in [H04] that for a domain Ω in a complex hermitian manifold of dimension n , if $\inf \sigma_e(\square_q) > 0$ for some q between 1 and $n-1$, then wherever the boundary is C^3 -smooth, its Levi-form cannot have exactly $n-q-1$ positive and q negative eigenvalues. A complete characterization of a domain in a complex hermitian manifold, in fact, even in $\mathbb{C}^n$, that has $\inf \sigma_e(\square_q) > 0$ or $\inf \sigma(\square_q) > 0$ is unknown.

Acknowledgments. We thank Professor Yum-Tong Siu for drawing our attention to the work of Laufer [L66], by which our work was inspired. We also thank Professors Mei-Chi Shaw and Emil Straube, and the referee for stimulating discussions and constructive suggestions.

REFERENCES

- [AG62] Aldo Andreotti and Hans Grauert, *Théorème de finitude pour la cohomologie des espaces complexes*, Bull. Soc. Math. France **90** (1962), 193-259.
- [BSt99] Harold P. Boas and Emil J. Straube, *Global regularity of the $\bar{\partial}$ -Neumann problem: a survey of the L^2 -Sobolev theory*, Several Complex Variables (M. Schneider and Y.-T. Siu, eds.), MSRI Publications, vol. 37, 79-112, 1999.
- [B02] Judith Brinkschulte, *Laufer's vanishing theorem for embedded CR manifolds*, Math. Z. **239**(2002), 863-866.

- [Br83] Thorsten Broecker, *Zur L^2 -Kohomologie beschränkter Gebiete*, Bonner Mathematische Schriften, vol. 145, Universitt Bonn, 1983.
- [C83] David Catlin, *Necessary conditions for subellipticity of the $\bar{\partial}$ -Neumann problem*, Ann. Math. **117** (1983), 147–171.
- [CS99] So-Chin Chen and Mei-Chi Shaw, *Partial differential equations in several complex variables*, AMS/IP, 2000.
- [Ch99] Michael Christ, *Remarks on global irregularity in the $\bar{\partial}$ -Neumann problem*, Several Complex Variables (M. Schneider and Y.-T. Siu, eds.), MSRI Publications, vol. 37, 161–198, 1999.
- [DK99] J. D’Angelo and J. J. Kohn, *Subelliptic estimates and finite type*, Several Complex Variables (M. Schneider and Y.-T. Siu, eds.), MSRI Publications, vol. 37, 199–232, 1999.
- [D95] E. B. Davies, *Spectral theory and differential operators*, Cambridge Studies in advanced mathematics, vol. 42, Cambridge University Press, 1995.
- [De] J.-P. Demailly, *Complex analytic and algebraic geometry*, book available at <http://www-fourier.ujf-grenoble.fr/~demailly/books.html>.
- [FK72] G. B. Folland and J. J. Kohn, *The Neumann problem for the Cauchy-Riemann complex*, Annals of Mathematics Studies, no. 75, Princeton University Press, 1972.
- [Fu05] Siqi Fu, *Hearing pseudoconvexity with the Kohn Laplacian*, Math. Ann. **331** (2005), 475–485.
- [Fu08] ———, *Hearing the type of a domain in C^2 with the d -bar-Neumann Laplacian*, Adv. in Math. **219** (2008), 568–603.
- [FS01] Siqi Fu and Emil J. Straube, *Compactness in the $\bar{\partial}$ -Neumann problem*, Complex Analysis and Geometry, Proceedings of Ohio State University Conference, vol. 9, 141–160, Walter De Gruyter, 2001.
- [GR65] Robert C. Gunning and Hugo Rossi, *Analytic functions of several complex variables*, Prentice-Hall, Englewood Cliffs, New Jersey, 1965.
- [H65] Lars Hörmander, *L^2 estimates and existence theorems for the $\bar{\partial}$ operator*, Acta Math. **113** (1965), 89–152.
- [H91] ———, *An introduction to complex analysis in several variables*, third ed., Elsevier Science Publishing, 1991.
- [H04] ———, *The null space of the $\bar{\partial}$ -Neumann operator*, Ann. Inst. Fourier (Grenoble) **54** (2004), 1305–1369.
- [HJ85] Roger A. Horn and Charles R. Johnson, *Matrix analysis*, Cambridge University Press, 1985.
- [Ka66] Marc Kac, *Can one hear the shape of a drum?* Amer. Math. Monthly **73** (1966), 1–23.
- [Ko63] J. J. Kohn, *Harmonic integrals on strongly pseudo-convex manifolds, I*, Ann. Math. **78** (1963), 112–148.
- [Ko64] ———, *Harmonic integrals on strongly pseudo-convex manifolds, II*, Ann. Math. **79** (1964), 450–472.
- [KN65] J. J. Kohn and L. Nirenberg, *Non-coercive boundary value problems*, Comm. Pure Appl. Math. **18** (1965), 443–492.
- [Kr01] Steven G. Krantz, *Function theory of several complex variables*, AMS Chelsea Publishing, Providence, RI, 2001.
- [L66] Henry B. Laufer, *On sheaf cohomology and envelopes of holomorphy*, Ann. Math. **84** (1966), 102–118.
- [L75] ———, *On the finite dimensionality of the Dolbeault cohomology groups*, Proc. Amer. Math. Soc. **52**(1975), 293–296.
- [O82] T. Ohsawa, *Isomorphism theorems for cohomology groups of weakly 1-complete manifolds*, Publ. Res. Inst. Math. Sci. **18** (1982), no. 1, 191–232.
- [O88] Takeo Ohsawa, *Complete Kähler manifolds and function theory of several complex variables*, Sugaku Expositions **1** (1988), 75–93.
- [Se53] J.-P. Serre, *Quelques problèmes globaux relatifs aux variétés de Stein*, Colloque sur les Fonctions de Plusieurs Variables, 57–68, Brussels, 1953.
- [Sh92] Mei-Chi Shaw, *Local existence theorems with estimates for $\bar{\partial}_b$ on weakly pseudo-convex CR manifolds*, Math. Ann. **294** (1992), no. 4, 677–700.
- [Sh09] ———, *The closed range property for $\bar{\partial}$ on domains with pseudoconcave boundary*, preprint, 2009, in this volume.
- [Siu67] Yum-Tong Siu, *Non-countable dimensions of cohomology groups of analytic sheaves and domains of holomorphy*, Math. Z. **102** (1967), 17–29.

- [Sk72] Henri Skoda, *Application des techniques L^2 à la théorie des idéaux d'une algèbre de fonctions holomorphes avec poids*, Ann. Sci. École Norm. Sup. (4) **5** (1972), 545-579.
- [St09] Emil Straube, *Lectures on the L^2 -Sobolev theory of the $\bar{\partial}$ -Neumann problem*, to appear in the series ESI Lectures in Mathematics and Physics, European Math. Soc.
- [W84] Jan Wiegerinck, *Domains with finite dimensional Bergman space*, Math. Z. **187** (1984), 559-562.

DEPARTMENT OF MATHEMATICAL SCIENCES, RUTGERS UNIVERSITY-CAMDEN, CAMDEN, NJ 08102
E-mail address: `sfu@camden.rutgers.edu`